

# ***Sphaerillo boninensis* Nunomura, 1990 (Crustacea, Isopoda, Oniscidea) is a junior synonym of a pantropical species, *Venezillo parvus* (Budde-Lund, 1885)**

Shigenori Karasawa<sup>1</sup>

<sup>1</sup> Department of Life and Environmental Agricultural Sciences, Faculty of Agriculture, Tottori University, 4-101 Koyama-machi Minami, Tottori 680-8553, Japan

Corresponding author: Shigenori Karasawa (dojyoudoubutu@gmail.com)

---

Academic editor: S. Taiti | Received 20 April 2018 | Accepted 14 February 2020 | Published 1 April 2020

---

<http://zoobank.org/B574D460-C84B-4D01-BD10-B4AB2D1C0C85>

---

**Citation:** Karasawa S (2020) *Sphaerillo boninensis* Nunomura, 1990 (Crustacea, Isopoda, Oniscidea) is a junior synonym of a pantropical species, *Venezillo parvus* (Budde-Lund, 1885). ZooKeys 923: 1–14. <https://doi.org/10.3897/zookeys.923.26018>

---

## **Abstract**

Re-examination of the holotype and paratype of *Sphaerillo boninensis* Nunomura, 1990 from Chichijima Island of the Ogasawara archipelago, which is registered as a UNESCO World Heritage Site, indicates that this species is a junior synonym of a pantropical species, *Venezillo parvus* (Budde-Lund, 1885).

## **Keywords**

Armadillidae, Chichijima Island, Ogasawara archipelago, terrestrial isopods, UNESCO World Heritage Site

## **Introduction**

*Sphaerillo boninensis* Nunomura, 1990, described from specimens collected on Chichijima Island, Ogasawara archipelago, is a small terrestrial isopod formerly considered endemic to this archipelago, a UNESCO World Heritage site approximately 1,000 km from the mainland of Japan. However, there are questions around the taxonomy of this species. The genus that this species has been attributed to has changed many times (see

the taxonomic account; Nunomura 1990, 1999a, 2011), and the species is included in the genus *Spherillo* in the current taxonomic treatment (Nunomura 2015). In addition, Saito (1993) recorded specimens collected from Chichijima Island and identified as the pantropical species *Venezillo parvus* (Budde-Lund, 1885) by Drs Ferrara and Taiti, but this species has been overlooked in reviews of Japanese terrestrial isopod species (Nunomura 1999a, b, 2015; Saito et al. 2000).

*Venezillo parvus*, originally and imperfectly described as *Armadillo parvus* by Budde-Lund (1885), was subsequently transferred to the genus *Venezillo* Verhoeff, 1928 (Green et al. 1990; Taiti and Ferrara 1991a). Today *Venezillo* accommodates more than 140 recognized species (Boyko et al. 2008). More comprehensive, subsequent descriptions of *V. parvus* provided more useful taxonomic characteristics than those present in the original description, i.e., in Schultz (1963) as *V. evergladensis* Schultz, 1963, a species now considered to be a junior synonym of *V. parvus*, Ferrara & Taiti, 1983 as *Sphaerillo* (?) *parvus* (Budde-Lund, 1885), Schmidt (2003) and Gregory (2014) as *V. parvus*. Figures provided by Schmidt (2003) depicted the basis of the male pereopod 7 as bearing dense setae on the anterior corner of the ventral side, a characteristic present also on the male pereopod 7 of *S. boninensis* (Nunomura 1990, fig. 146I). Additionally, the oblique lobe on the ventral surface of pereonite 2 reported for *S. boninensis* by Nunomura (1990, fig. 146B) is a diagnostic feature of the genus *Venezillo* (Verhoeff 1928; Vandel 1952). These similarities indicated that *S. boninensis* was referable to *Venezillo* and a close relationship between *S. boninensis* and *V. parvus* might exist.

Terrestrial isopods are now scarce in the southern part of Hahajima Island, Ogasawara archipelago, in an area invaded by the land nemertean *Geonemertes pelaensis* Semper, 1863 (Shinobe et al. 2017). This terrestrial predator poses a threat to native biodiversity in this region. Resolving the systematic status of *S. boninensis*, or at least specimens referred to it, from this area would improve our understanding of actual versus perceived threats to endemic faunas of this region, and ultimately, the conservation and management of species throughout this archipelago.

The aim of this study was to resolve relationships between *S. boninensis* and *V. parvus*. This is achieved through re-examination and redescription of appropriate type material of the former.

## Materials and methods

### Sample collection

Type material was loaned from the Toyama Science Museum (collection acronym TOYA). As the previously dissected male holotype and paratype of *S. boninensis* were in poor condition, a description of the whole body, cephalon, antenna 1, and pleopod 5 of the female paratype was necessary. Additionally, as the male pereopods 1 and 7 are broken in male holotype, they can be only partly described. Male pereopod 2 was not found in the holotype and paratype.

## Morphology observation

Antennae 1 and 2, mandible, maxillae 1 and 2, maxilliped, pleopods, and pereopods were placed in Hoyer's mounting medium (Krantz and Walter 2009) on slides, covered with a coverslip, and drawn under a microscope (magnification of 40–400 ×; Eclipse E400, NIKON Corp.). The whole body, cephalon, and epimera of pereonites 1–7 were examined using a Nikon SMZ1500 microscope (magnification of 7.5–112.5 ×).

## Results

### Taxonomic account

#### Genus *Venezillo* Verhoeff, 1928

##### *Venezillo parvus* (Budde-Lund, 1885)

*Armadillo parvus* Budde-Lund, 1885: 25–26; Dollfus 1893: 187; Verhoeff 1946: 4; Jeppesen 2000: 254.

*Spherillo parvus*: Budde-Lund 1904: 91; Budde-Lund 1908: 270–271, taf 12, figs 30–37; Budde-Lund 1912: 371; Barnard 1964: 53.

*Venezillo evergladensis* Schultz, 1963: 209–213, figs 1–26; Schultz 1972: 1–4, figs 4–6; Schultz 1975: 186; Johnson 1976: 157, fig. 1; Johnson 1977: 603; Johnson 1978: 140; Johnson 1980: 124; Johnson 1981: 351, fig. 1; Johnson 1982: 225; Johnson 1983: 209; Johnson 1984: 465; Johnson 1985a: 403; Johnson 1985b: 216; Keeney 1990: 1–2, figs 1, 2.

*Sphaerillo parvus*: Ferrara and Taiti 1979: 182.

*Sphaerillo* (?) *parvus*: Ferrara and Taiti 1981: 196; Ferrara and Taiti 1983: 70–71, figs 131–136; Taiti and Ferrara 1983: 220–222.

*Sphaerillo boninensis* Nunomura, 1990: 19–21, fig. 146. Syn. nov.

*Venezillo parvus*: Green et al. 1990: 431, 433; Taiti and Ferrara 1991a: 220; Taiti and Ferrara 1991b: 915; Kwon and Taiti 1993: 77; Kwon 1995: 533; Taiti and Howarth 1996: 68; Taiti et al. 1998: 300; Leistikow and Wägele 1999: 49; Taiti 1999: 38; Jass and Klausmeier 2000: 781; Stoyenoff 2001: tab. 2; Green et al. 2002: 301; Zimmer et al. 2002: 597; Buck et al. 2003: 106; Kelly and Samways 2003: app 1; Schmidt 2003: 107, figs 122–128; Soesbergen 2003: 97, figs 1, 2, tab. 1; Schmalfuss 2004: 291; Zimmer et al. 2004: 754, figs 1–3, 5–12; Dias et al. 2005: tab. 3; McLaughlin et al. 2005: 198; Poore 2005: 11; Berg et al. 2008: 61; Gregory 2009a: 7; Gregory 2009b: 1; Barber 2010: 73; Cochard et al. 2010: tab. 7.1.1; Senterre et al. 2011: app 2; Murphy et al. 2012: 3; Arab and Wimp 2013: 333; Treplin et al. 2013: 84; Wimp et al. 2013: 507, fig. 1; Angelini and Siliman 2014: 188; Gregory 2014: 21–23, fig. 11, tab. 3; Humphreys 2014: tab. 2; Lemaire and Gerriet 2014: 43–44; Taiti and Wynne 2015: 44; Taiti 2018: 88;

Vittori and Štrus 2017: 396; De los Rios-Escalante et al. 2018: 50, tab. 1; Pérez-Schultheiss et al. 2018: 8.


*Venezillo boninensis*: Nunomura 1999a: 89; Nunomura 1999b: 598, 614, 624; Saito et al. 2000: 94. Syn. nov.

*Sphaerillo boninensis*: Nunomura 2011: 71; Nunomura 2015: 1032, 1050, 1063; Shinobe et al. 2017: 3. Syn. nov.


**Material examined.** *Sphaerillo boninensis*: Holotype, TOYA-Cr-8953, male, dissected, forest of *Casuarina equisetifolia*, Suzaki, Chichijima Island, Tokyo Metropolis, Japan, 1 July 1977, Jun-Ichi Aoki leg; paratype, TOYA-Cr-8955, male, dissected, forest of *Casuarina equisetifolia*, Suzaki, Chichijima Island, Tokyo Metropolis, Japan, 1 July 1977, Jun-Ichi Aoki leg; paratype, TOYA-Cr-8961, female, dissected, forest of *Casuarina equisetifolia*, Suzaki, Chichijima Island, Tokyo Metropolis, Japan, 1 July 1977, Jun-Ichi Aoki leg.

### Redescription of *Sphaerillo boninensis* Nunomura, 1990

Body color yellowish in alcohol. Pereonites 1–7 with single nodulus lateralis per side, all similarly distanced from lateral margin (Fig. 1A). Pereonite 1 with lateral margin not grooved; schisma deep, with rounded inner and outer lobes, outer lobe protruding posteriorly compared to inner lobe (Fig. 1A, B). Pereonite 2 with an oblique lobe on ventral surface (Fig. 1C). Pereonite 3 without ventral lobe (Fig. 1D). Pereonites 4–7 with a small lobe on ventral surface (Fig. 1D–H). Eyes (of female paratype) with ten ommatidia (Fig. 1I, J). Upper middle edge of cephalon convex; frontal shield separated from vertex, trapezoidal in frontal view (Fig. 1I, J). Flagellum of second antenna with two articles (Fig. 1K). First antenna of three articles; apical article with numerous aesthetascs (Fig. 1L). Right mandible with two plumose setae between lacinia mobilis and molar penicil; lacinia mobilis of left mandible larger than right mandible; molar penicil unbranched; left mandible with three plumose setae between hairy lobe and molar penicil (Fig. 2A, B). First maxilla outer endite with ten simple teeth; inner endite with two stout plumose penicils (Fig. 2C, D). Second maxilla apically bilobate, covered with short setae (Fig. 2E). Endite of maxilliped rectangular, bearing three spines on distal margin; maxilliped palp with basal article bearing two long setae, distal article with apical tuft of small setae (Fig. 2F). Male pereopod 1 with antennal brush on carpus; propodus with numerous short setae on basal half of inner margin; carpus with five long stout setae on inner margin; apical point of outer margin of merus with two tapering setae, inner margin with three branched setae (Fig. 3A). Male pereopods 3–6 unremarkable (Fig. 3B–E). Basis of male pereopod 7 with dense setae on apical corner of inner margin (Fig. 3F). All pleopod exopodites with monospiracular covered lungs (Fig. 4). Male pleopod 1 with straight endopodite, with apical part bent outward with long setae; inner margin with row of small setae; outer margin with at least three tooth-like setae (Fig. 4A); exopodite triangular, with small setae on inner margin (Fig. 4B). Male pleopod 2 endopodite slender (apical part



**Figure 1.** *Sphaerillo boninensis* **A** whole body, lateral view, paratype (female) **B-H** epimeron of pleonites 1-7, ventral view, holotype (male) **I** cephalon, frontal view, paratype (female) **J** cephalon, dorsal view, paratype (female) **K** second antenna, holotype (male) **L** first antenna, paratype (female). Scale bars: 1.5 mm (**A**); 100 µm (**B-H, K**); 300 µm (**I, J**); 50 µm (**L**).




**Figure 2.** *Sphaerillo boninensis* **A** right mandible, holotype (male) **B** left mandible, holotype (male) **C** outer endite of first maxilla, holotype (male) **D** inner endite of first maxilla, holotype (male) **E** second maxilla, holotype (male) **F** maxilliped, holotype (male). Scale bars: 100 µm (**A, B**); 50 µm (**C–F**).

broken); exopodite triangular, with distal half elongated (Fig. 4C). Male pleopod 3 exopodite with triangular part on posterior inner corner (Fig. 4D). Male pleopod 4 exopodite parallelogram-shaped (Fig. 4E). Female pleopod 5 parallelogram-shaped, with fine setae of variable length along inner margin (Fig. 4F). Pleotelson hour-glass-shaped, distal part narrower than basal part (Fig. 5A). Uropodal protopod trapezoidal, with exopodite inserted on dorsal surface of protopod; endopodite about twice as long as exopodite (Fig. 5B).


## Discussion

The genus *Venezillo* is characterized among other characters by a narrow lobe located obliquely or horizontally on the ventral surface of the pereon epimeron 2 (Verhoeff




**Figure 3.** *Sphaerillo boninensis* **A** pereopod 1, holotype (male) **B-E** pereopods 3–6, holotype (male) **F** basis of pereopod 7, holotype (male). Scale bar: 100 µm.

1928; Vandel 1952). The holotype of *S. boninensis* has an oblique lobe on the ventral surface of the epimeron 2, for which reason it is more appropriately assigned to the genus *Venezillo*. In addition, in all other morphological characteristics, type materials of *S. boninensis* are consistent in morphology with those of *V. parvus* as redescribed by Schmidt (2003): the apical corner of the basis of the male pereopod 7 bears dense short setae (Fig. 3F), the endopodite of the male pleopod 1 has long setae at the tip and scale-like setae on the inner margin (Fig. 4A), and the exopodite of the male pleopod 1



**Figure 4.** *Sphaerillo boninensis* **A** endopodite of pleopod 1, holotype (male) **B** exopodite of pleopod 1, holotype (male) **C–E** pleopods 2–4, holotype (male) **F** pleopod 5, paratype (female). Scale bar: 100 µm.



**Figure 5.** *Sphaerillo boninensis* **A** pleotelson, holotype (male) **B** uropod, holotype (male). Scale bars: 200 µm (**A**); 100 µm (**B**).

is triangular (Fig. 4B). For these reasons, I regard *S. boninensis* to be a junior synonym of *V. parvus*, a species widely distributed in the tropical belt.

While *S. boninensis* was considered endemic to the Ogasawara archipelago, and potentially threatened by an invasive nemertean predator (Shinobe et al. 2017), the pantropical *V. parvus*, due to its widespread distribution and invasive tendency, is not considered to be threatened at species level.

## Acknowledgements

I would like to thank Dr Tomofumi Iwata (Toyama Science Museum) and Mr Noboru Nunomura (Kanazawa University, Institute of Nature and Environmental Technology) for loaning type materials of *Sphaerillo boninensis*. This work was supported by a Grant-in-Aid for Young Scientists (B) Grant Number 26830145.

## References

- Angelini C, Silliman BR (2014) Secondary foundation species as drivers of trophic and functional diversity: evidence from a tree-epiphyte system. *Ecology* 95: 185–196. <https://doi.org/10.1890/13-0496.1>
- Arab A, Wimp GM (2013) Plant production and alternate prey channels impact the abundance of top predators. *Oecologia* 173: 331–341. <https://doi.org/10.1007/s00442-013-2618-7>
- Barber T, Gregory S, Lee P (2010) Reports on the 2009 BMIG spring meeting in Cornwall. *Bulletin of the British Myriapod Isopod Group* 24: 65–74.
- Barnard KH (1964) The terrestrial Isopoda and Amphipoda of the Mascarene Islands. *Bulletin of the Mauritius Institute* 6: 49–60.

- Berg MP, Soesbergen M, Tempelman D, Wijnhoven H (2008) Verspreidingsatlas Nederlandse landpissebedden, Duizendpoten en Miljoenpoten (Isopoda, Chilopoda, Diplopoda). EIS-Nederland, Leiden Vrije Universiteit-Afdeling Dierecologie, Amsterdam, 194 pp.
- Boyko CB, Bruce NL, Hadfield KA, Merrin KL, Ota Y, Poore GCB, Taiti S, Schotte M, Wilson GDF (2008) World Marine, Freshwater and Terrestrial Isopod Crustaceans database. *Venezillo Verhoeff*, 1928. Accessed through: World Register of Marine Species. <http://www.marinespecies.org/aphia.php?p=taxdetails&id=249464> on 2019-03-15
- Buck TL, Breed GA, Pennings SC, Chase ME, Zimmer M, Carefoot TH (2003) Diet choice in an omnivorous salt-marsh crab: different food types, body size, and habitat complexity. *Journal of Experimental Marine Biology and Ecology* 292: 103–116. [https://doi.org/10.1016/S0022-0981\(03\)00146-1](https://doi.org/10.1016/S0022-0981(03)00146-1)
- Budde-Lund G (1885) Crustacea Isopoda Terrestria, per Familias et Genera et Species Descripta. Hauniæ, Copenhagen, 319 pp. <https://doi.org/10.5962/bhl.title.109769>
- Budde-Lund G (1904) Revision of Crustacea Isopoda Terrestria, with Additions and Illustrations. pt. 1. *Eubelum*, pt. 2 Spherilloniinae, pt. 3 *Armadillo*. H. Hagerup, Copenhagen. <https://doi.org/10.5962/bhl.title.9883>
- Budde-Lund G (1908) Isopoda von Madagaskar und Ostafrika. Mit diagnosen verwandter arten. Reise in Ostafrika in den Jahren 1903-1905, Wissenschaftliche Ergebnisse 2: 265–308. <https://doi.org/10.5962/bhl.title.12989>
- Budde-Lund G (1912) Terrestrial Isopoda, Particularly Considered in Relation to the Distribution of the Southern Indopacific Species. *Transactions of the Linnean Society of London*, 2<sup>nd</sup> Series, Zoology. <https://doi.org/10.1111/j.1096-3642.1912.tb00107.x>
- Cochard PO, Vilisics F, Sechet E (2010) Alien terrestrial crustaceans (Isopods and Amphipods). Chapter 7.1. In: Roques A, Kenis M, Lees D, Lopez-Vaamonde C, Rabitsch W, Rasplus J-Y, Roy D (Eds) Alien terrestrial arthropods of Europe. *BioRisk* 4: 81–96. <https://doi.org/10.3897/biorisk.4.54>
- De los Rios-Escalante P, Arancibia EI, Pérez-Schultheiss J (2018) A checklist of non-marine crustaceans from Chilean oceanic islands. *Proceedings of the Biological Society of Washington* 131: 47–52. <https://doi.org/10.2988/17-00007>
- Dias N, Sprung M, Hassall M (2005) The abundance and life histories of terrestrial isopods in a salt marsh of the Ria Formosa lagoon system, southern Portugal. *Marine Biology* 147: 1343–1352. <https://doi.org/10.1007/s00227-005-0033-2>
- Dollfus A (1893) Voyage de M. Charles Alluaud aux Iles Sécherelles. Crustacés isopodes terrestres. *Bulletin de la Société Zoologique de France* 18: 186–190.
- Ferrara F, Taiti S (1979) A check-list of terrestrial isopods from Africa (south of the Sahara). *Monitore Zoologico Italiano* (Nuova Serie) Supplemento 12: 89–215. <https://doi.org/10.1080/03749444.1979.10736595>
- Ferrara F, Taiti S (1981) Terrestrial isopods from Ascension Islands. *Monitore Zoologico Italiano* (Nuova Serie) Supplemento 14: 189–198. <https://doi.org/10.1080/03749444.1981.10736621>
- Ferrara F, Taiti S (1983) Isopodi terrestri: contributions à l'étude de la faune terrestre des îles granitiques de l'archipel des Sécherelles (Mission P.L.G. Benoit, J.J. Van Mol 1972). *Annales Du Musée Royal De l'Afrique Centrale*, Série 8vo, Sciences Zoologiques 240: 1–92.

- Green AJA, Ferrara F, Taiti S (1990) Terrestrial isopoda from the Krakatau islands, south Sumatra and west Java. Memoirs of the Museum of Victoria 50: 417–436. <https://doi.org/10.24199/j.mmv.1990.50.13>
- Green AJA, Lew Ton HM, Poore GCB (2002) Suborder: Oniscidea Latreille, 1802. In: Poore GCB (Ed.) Zoological Catalogue of Australia 19.2A: Crustacea: Malacostraca: Syncarida, Peracarida, Isopoda, Tanaidacea, Mictacea, Thermosbaenacea, Spelaeogriphacea. CSIRO Publishing, Melbourne, 279–344.
- Gregory S (2009a) Woodlice and Waterlice (Isopoda: Oniscidea & Asellota) in Britain and Ireland. FSC Publications, Shrewsbury, 175 pp.
- Gregory S (2009b) A tropical woodlouse new to Britain. British Myriapod and Isopod Group Newsletter 18: 1–2.
- Gregory S (2014) Woodlice (Isopoda: Oniscidea) from the Eden Project, Cornwall, with descriptions of new and poorly known British species. Bulletin of the British Myriapod Isopod Group 27: 3–27.
- Humphreys WF (2014) Subterranean fauna of Christmas Island: habitats and salient features. Raffles Bulletin of Zoology Supplement 30: 29–44.
- Jass J, Klausmeier B (2000) Endemics and immigrants: North American terrestrial isopods (Isopoda, Oniscidea) north of Mexico. Crustaceana 73: 771–799. <https://doi.org/10.1163/156854000504804>
- Jeppesen PC (2000) Catalogue of terrestrial isopod taxa and type material described by Gustav Budde-Lund (Crustacea: Isopoda). Steenstrupia 25: 221–265.
- Johnson C (1976) Genetics of red body polymorphism in the isopod, *Venezillo evergladensis*. Journal of Heredity 67: 157–160. <https://doi.org/10.1093/oxfordjournals.jhered.a108693>
- Johnson C (1977) Evolution of sex ratios in the isopod, *Venezillo evergladensis*. Evolution 31: 603–610. <https://doi.org/10.1111/j.1558-5646.1977.tb01049.x>
- Johnson C (1978) A note on codominance in the terrestrial isopod, *Venezillo evergladensis*. Journal of Heredity 69: 1–140. <https://doi.org/10.1093/oxfordjournals.jhered.a108902>
- Johnson C (1980) Genetics of albinism in the isopod. Journal of Heredity 71: 124–126. <https://doi.org/10.1093/oxfordjournals.jhered.a109323>
- Johnson C (1981) Genetics of mottling in the terrestrial isopod. Journal of Heredity 72: 351–352. <https://doi.org/10.1093/oxfordjournals.jhered.a109521>
- Johnson C (1982) Multiple insemination and sperm storage in the isopod, *Venezillo evergladensis* Schultz, 1963. Crustaceana 42: 225–232. <https://doi.org/10.1163/156854082X00920>
- Johnson C (1983) Eye color genetics and allelism of r-locus factors in the Isopod, *Venezillo evergladensis*. Journal of Heredity 74: 209–210. <https://doi.org/10.1093/oxfordjournals.jhered.a109770>
- Johnson C (1984) Geographic variability in the genetic polymorphism of the isopod, *Venezillo evergladensis*. Journal of Heredity 75: 465–474. <https://doi.org/10.1080/00222938400770391>
- Johnson C (1985a) Selection dynamics in the isopod *Venezillo evergladensis*. Pedobiologia 28: 403–412.
- Johnson C (1985b) Mating behavior of the terrestrial isopod, *Venezillo evergladensis* (Oniscidea, Armadillidae). The American Midland Naturalist 114: 216–224. <https://doi.org/10.2307/2425597>

- Kelly JA, Samways MJ (2003) Diversity and conservation of forest-floor arthropods on a small Seychelles island. *Biodiversity and Conservation* 12: 1793–1813. <https://doi.org/10.1023/A:1024161722449>
- Keeney GD (1990) Some exotic terrestrial isopods (Oniscoidea) from the Columbus Zoo Exploration Center, Powell, Ohio: two new state records. *Ohio Journal of Science* 90: 133–134. <http://hdl.handle.net/1811/23407>
- Krantz W, Walter E (2009) A Manual of Acarology (3<sup>rd</sup> ed). Texas Tech University Press, Texas, 807 pp.
- Kwon DH (1995) Terrestrial Isopoda (Crustacea) from Cheju Island, Korea. *The Korean Journal Systematic Zoology* 11: 509–538.
- Kwon DH, Taiti S (1993) Terrestrial Isopoda (Crustacea) from southern China, Macau and Hong Kong. *Stuutgarter Beitr Naturk (Serie a)* 490: 1–83. <https://www.biodiversitylibrary.org/page/33439218#page/79/mode/1up>
- Leistikow A, Wägele JW (1999) Checklist of the terrestrial isopods of the new world (Crustacea, Isopoda, Oniscidea). *Revista Brasileira de Zoologia* 16: 1–72. <https://doi.org/10.1590/S0101-81751999000100001>
- Lemaire JM, Gerriet O (2014) Les invertébrés (Arthropoda & Mollusca) de la grande serre du Parc Phoenix (Nice, Alpes-Maritimes, France). *Riviera Scientifique* 98: 39–52.
- McLaughlin PA, Camp DK, Angel MV, Bousfi EL, Brunel P, Brusca RC, Cadien D, Cohen AC, Conlan K, Eldredge LG, Felder DL, Goy JW, Haney R, Haan B, Heard RW, Hendrycks EA, Hobbs III HH, Holsinger JR, Kensley B, Laubitz DR, LeCroy SE, Lemaitre R, Maddocks RF, Martin JW, Mikkelsen P, Nelson E, Newman WA, Overstreet RM, Poly WJ, Price WW, Reid JW, Robertson A, Rogers DC, Ross A, Schram FR, Shih CT, Watling L, Wilson GDF, Turgeon DD (2005) Common and scientific names of aquatic invertebrates from the United States and Canada: crustaceans. *American Fisheries Society Special Publication* 31: 1–533.
- Murphy SM, Wimp GM, Lewis D, Denno RF (2012) Nutrient presses and pulses differentially impact plants, herbivores, detritivores and their natural enemies. *PLoS ONE* 7: e43929. <https://doi.org/10.1371/journal.pone.0043929>
- Nunomura N (1990) Studies on the terrestrial isopod crustaceans in Japan V. Taxonomy of the families Armadillidiidae, Armadillidae and Tylidae, with taxonomic supplements to some other families. *Bulletin of the Toyama Science Museum* 13: 1–58.
- Nunomura N (1999a) Taxonomical revisions on some groups of terrestrial isopods in Japan. *Edaphologia* 62: 81–91. [In Japanese]
- Nunomura N (1999b) Crustacea, Isopoda. In: Aoki J (Ed.) *Pictorial Keys to Soil Animals of Japan*. Tokai University Press, Tokyo, 569–625. [In Japanese]
- Nunomura N (2011) Crustaceans No.2 (Isopoda). *Special Publication of the Toyama Science Museum* 24: 1–133. [In Japanese]
- Nunomura N (2015) Crustacea, Isopoda. In: Aoki J (Ed.) *Pictorial Keys to Soil Animals of Japan*. Tokai University Press, Kanagawa, 997–1065. [In Japanese]
- Pérez-Schultheiss J, Ayala K, Fariña JM, Coccia C (2018) Exotic oniscideans (Crustacea: Isopoda) in coastal salt marshes: first record of the families Halophilosciidae and Platyarthridae in continental Chile. *New Zealand Journal of Zoology*: 1–11. <https://doi.org/10.1080/03014223.2018.1539017>

- Poore GCB (2005) Supplement to the 2002 catalogue of Australian Crustacea: Malacostraca—Syncarida and Peracarida (Volume 19.2A): 2002–2004. Museum Victoria Science Reports 7: 1–15. <https://doi.org/10.24199/j.mvsr.2005.07>
- Saito S (1993) Terrestrial crustaceans. In: Society of Nature Study of Yanba Dam (Ed.) *Nature of Naganohara*. Naganohara Town, Gunma, 345–356. [In Japanese]
- Saito N, Itani G, Nunomura N (2000) A preliminary check list of isopod crustaceans in Japan. Bulletin of the Toyama Science Museum 23: 11–107. [In Japanese]
- Schmalfuss H (2004) World catalog of terrestrial isopods (Isopoda: Oniscidea). <http://www.oniscidea-catalog.naturkundemuseum-bw.de>
- Schmidt C (2003) Contribution to the phylogenetic system of the Crinocheta (Crustacea, Isopoda). Part 2. (Oniscoidea to Armadillidiidae). Mitteilungen aus dem Museum für Naturkunde in Berlin, Zoologische Reihe 79: 3–179. <https://doi.org/10.1002/mmzn.20030790102>
- Schultz GA (1963) *Venezillo evergladensis*, a new species of terrestrial isopod crustacean from Florida. Transactions of the American Microscopical Society 82: 209–213. <https://doi.org/10.2307/3223996>
- Schultz GA (1972) The Armadillidae of Florida (Isopoda, Oniscoidea). Quarterly Journal of the Florida Academy of Sciences 35: 1–4. <https://www.jstor.org/stable/24318279>
- Schultz GA (1975) Terrestrial isopod crustaceans (Oniscoidea) from coastal sites in Georgia. Bulletin of the Georgia Academy of Science 34: 185–194.
- Senterre B, Henriette E, Chong-Seng L, Beaver K, Mougal J, Vel T, Gerlach J (2011) Seychelles Key Biodiversity Areas—Output 1: List of species of special concern. Consultancy Report, Ministry of Environment-UNDP-GEF Project, Victoria, 67 pp.
- Shinobe S, Uchida S, Mori, H, Okochi I, Chiba S (2017) Declining soil Crustacea in a World Heritage Site caused by land nemertean. Scientific Reports 7: 1–8. <https://doi.org/10.1038/s41598-017-12653-4>
- Soesbergen M (2003) *Venezillo parvus* en *Synarmadillo* spec., twee nieuwe land-pissebedden in Nederland (Crustacea: Isopoda: Armadillidae). Nederlandse Faunistische Mededelingen 18: 97–101.
- Stoyenoff JL (2001) Distribution of terrestrial isopods (Crustacea: Isopoda) throughout Michigan: early results. The Great Lakes Entomologist 34: 29–50. <https://scholar.valpo.edu/tgle/vol34/iss2/4>
- Taiti S (1999) Terrestrial isopods from the Midway Islands. Occasional Papers of the Bishop Museum 58: 37–38.
- Taiti S (2018) Biology and biogeography of terrestrial isopods (Crustacea, Isopoda, Oniscidea). Atti Accademia Nazionale Italiana di Entomologia 65: 83–90.
- Taiti S, Ferrara F (1983) Su alcuni isopodi terrestri della Réunion, di Mauritius e delle Seycelles. Revue Suisse de Zoologie 90: 199–231. <https://doi.org/10.5962/bhl.part.81973>
- Taiti S, Ferrara F (1991a) Terrestrial isopods (Crustacea) from the Hawaiian Islands. Bishop Museum Occasional Papers 31: 202–227.
- Taiti S, Ferrara F (1991b) Two new species of terrestrial Isopoda (Crustacea, Oniscidea) from Ascension Island. Journal of Natural History 25: 901–916. <https://doi.org/10.1080/00222939100770591>

- Taiti S, Howarth FG (1996) Terrestrial isopods from the Hawaiian Islands (Isopoda: Oniscidea). Bishop Museum Occasional Papers 45: 59–71. <http://hbs.bishopmuseum.org/pdf/op45-59-71.pdf>
- Taiti S, Paoli P, Ferrara F (1998) Morphology, biogeography, and ecology of the family Armadillidae (Crustacea, Oniscidea). Israel Journal of Zoology 44: 291–301. <https://www.tandfonline.com/doi/abs/10.1080/00212210.1998.10688952>
- Taiti S, Wynne JJ (2015) The terrestrial Isopoda (Crustacea, Oniscidea) of Rapa Nui (Easter Island), with descriptions of two new species. In: Taiti S, Hornung E, Štrus J, Bouchon D (Eds) Trends in Terrestrial Isopod Biology. ZooKeys 515: 27–49. <https://doi.org/10.3897/zookeys.515.9477>
- Treplin M, Pennings SC, Zimmer M (2013) Decomposition of leaf litter in a U.S. saltmarsh is driven by dominant species, not species complementarity. Wetlands 33: 83–89. <https://doi.org/10.1007/s13157-012-0353-1>
- Vandel A (1952) Etude des isopodes terrestres récoltés au Vénézuela par le Dr. G. Marcuzzi, suivie de considerations sur le peuplement du Continent de Gondwana. Memoire del Museo Civico di Storia Naturale di Verona 3: 59–203.
- Verhoeff KW (1928) Über einige Isopoden der zoologischen Staatssammlung in München. 38. Isopoden-Aufsatz. Zoologischer Anzeiger 76: 113–123.
- Verhoeff KW (1946) Über Land-Isopoden der Seychellen und aus Burma. Arkiv för Zoologi 37: 1–18.
- Vittori M, Štrus J (2017) Male tegumental glands in the cavernicolous woodlouse *Cyphonethes herzegowinensis* (Verhoeff, 1900) (Crustacea: Isopoda: Trichoniscidae). Journal of Crustacean Biology 37: 389–397. <https://doi.org/10.1093/jcobi/rux067>
- Wimp GM, Murphy SM, Lewis D, Douglas MR, Ambikapathi R, Van-Tull L, Gratton C, Denno RF (2013) Predator hunting mode influences patterns of prey use from grazing and epigaeic food webs. Oecologia 171: 505–515. <https://doi.org/10.1007/s00442-012-2435-4>
- Zimmer M, Pennings SC, Buck TL, Carefoot TH (2002) Species-specific patterns of litter processing by terrestrial isopods (Isopoda: Oniscidea) in high intertidal salt marshes and coastal forests. Functional Ecology 16: 596–607. <https://doi.org/10.1046/j.1365-2435.2002.00669.x>
- Zimmer M, Pennings SC, Buck TL, Carefoot TH (2004) Salt marsh litter and detritivores: A closer look at redundancy. Estuaries 27: 753–769. <https://doi.org/10.1007/BF02912038>